

Saturated magnetic field amplification at supernova shocks

Qinghuan Luo and Don Melrose

School of Physics, The University of Sydney, NSW 2006, Australia

— Received in original form February, 2009

ABSTRACT

Cosmic-ray streaming instabilities at supernova shocks are discussed in the quasilinear diffusion formalism which takes into account the feedback effect of wave growth on the cosmic ray streaming motion. In particular, the nonresonant instability that leads to magnetic field amplification in the short wavelength regime is considered. The linear growth rate is calculated using kinetic theory for a streaming distribution. We show that the nonresonant instability is actually driven by a compensating current in the background plasma. The nonresonant instability can develop into a nonlinear regime generating turbulence. The saturation of the amplified magnetic fields due to particle diffusion in the turbulence is derived analytically. It is shown that the evolution of parallel and perpendicular cosmic-ray pressures is predominantly determined by non-resonant diffusion. However, the saturation is determined by resonant diffusion which tends to reduce the streaming motion through pitch angle scattering. The saturated level can exceed the mean background magnetic field.

Key words: ISM: magnetic field – cosmic rays – supernova remnants – shock – radiation mechanisms

1 INTRODUCTION

Diffusive shock acceleration (DSA) is regarded as the preferred mechanism for the acceleration of the Galactic cosmic rays (CRs) (Drury 1983). It is commonly believed that SN shocks are responsible for acceleration of high energy CRs at least up to the ‘knee’ ($\sim 4 \times 10^{15}$ eV) of the CR spectrum (Hillas 2006). In the DSA model, particles gain energy by bouncing back and forth across the shock. Although particles only gain a small amount of energy in each crossing, they can be accelerated to very high energy through many crossings provided they can be trapped in the acceleration region for a sufficiently long time. The acceleration is rather efficient and naturally leads to a power-law energy distribution. Despite these advantages, there have been two long standing problems. First, the standard DSA theory predicts the maximum energy well below the ‘knee’ (Bell 2004). The maximum energy achievable is limited by both the Bohm approximation, in which the particle’s gyroradius must not exceed the mean free path to scattering, and the condition that the gyroradius is smaller than the shock’s width (otherwise the particle would escape from the acceleration region) (Bell 2004; Zirakashvili, Ptuskin & Völk 2008). For a magnetic field of order 10^{-10} T and typical parameters of SN shocks, one has the maximum energy $\sim 5 \times 10^{14}$ eV. Second, turbulence is required for effective scattering in the acceleration region so that the particles can be trapped (Skilling 1975; Bell 1978; Lagage & Cesarsky 1983a,b). So far, in the standard DSA theory, one generally postulates that Alfvén

turbulence can be generated by CRs themselves through resonant interactions, even though in practice the growth of Alfvén waves due to CRs is known to be ineffective.

There is growing interest in the possibility that the magnetic field at the shock may be amplified due to CR induced instability in the nonresonant regime (McKenzie & Völk 1982; Bell & Lucek 2001; Bell 2004). Bell (2004) showed a nonresonant form of the instability driven by a CR current can outgrow the familiar resonant form, leading to magnetic amplification. A strong magnetic field at shocks reduces the gyroradius and this raises the maximum energy to which particles can be accelerated. X-ray observations of young SNRs suggest that the magnetic field strength near the shocks is much higher than that in the ISM and is a strong function of the shock speed (Vink & Laming 2003; Völk, Ksenofontov & Berezhko 2008). The presence of strong magnetic fields may explain the lack of strong TeV gamma-ray fluxes, as shown from HESS observations (Völk, Ksenofontov & Berezhko 2008). The nonthermal X-rays are well described by synchrotron spectra. A stronger magnetic field in the emission region implies that a lower electron energy is required. This would lead to a lower TeV gamma flux from inverse Compton scattering.

The nonresonant growth of Alfvén waves due to CR streaming has been discussed in both the MHD formalism (Bell & Lucek 2001; Bell 2004; Zirakashvili, Ptuskin & Völk 2008) and the kinetic formalism (Melrose 2005; Reville, Kirk & Duffy 2006; Amato & Blasi 2008). In both formalisms, the instability is shown to exist in the

linear regime. To test the magnetic field amplification model against observations one needs to determine the saturated magnetic field accurately. The saturation of the instability cannot be determined in linear theory as the reaction of the instability on the CR momentum distribution is not automatically included in the linear calculation. Although there are numerical simulations of the instability that extend to the nonlinear regime, different saturation levels have been predicted (Niemi, Pohl, Stroman, & Nishikawa 2008; Riquelme & Spitkovsky 2008). In this paper, the instability is discussed in the quasilinear formalism in which the reaction of the instability on the CR distribution can be included self-consistently. So, in this formalism one can estimate the saturation analytically, with both nonresonant and resonant diffusion processes considered. The treatment of the nonresonant diffusion presented here is similar to that used for the firehose instability (Davidson 1972). We emphasize the major difference between the CR streaming instability and the firehose instability. The former is caused by streaming motion and the latter is due to a pressure anisotropy with excess of parallel pressure over the perpendicular pressure (with respect to the mean magnetic field). To some extent, the streaming instability also resembles the Weibel instability—a nonresonant, purely growing mode driven by anisotropy in the particle distribution (Weibel 1959).

In Sec 2 the kinetic theory of CR streaming instabilities is discussed with emphasis on the nonresonant instability. Quasilinear diffusion driven by the nonresonant instability is discussed in both the short and long wavelength approximations in the nonresonant regime in Sec 3 and in the resonant regime in Sec 4. Application to SN shocks is discussed in Sec 5.

2 COSMIC-RAY STREAMING INSTABILITIES

We outline the kinetic theory of CR streaming instabilities including both the usual resonant instability and the nonresonant instability and focus particularly on the latter. Our treatment builds on other recent discussions of linear kinetic theory of the CR-induced nonresonant instability (Reville, Kirk & Duffy 2006; Amato & Blasi 2008). For convenience we assume a single species of CRs with charge q and mass m .

2.1 CR streaming motion

To model CR streaming at velocity v_{CR} along the mean magnetic field we consider a class of streaming distributions in momentum space given by

$$f(u_{\parallel}, u_{\perp}) = n_{CR} \left[1 + (2\sigma + 1) \frac{v_{CR}}{v} \left(\frac{u_{\parallel}}{u} \right)^{2\sigma-1} \right] \frac{g(u)}{4\pi u^2}, \quad (1)$$

where $u_{\parallel}$ and $u_{\perp}$ are the nondimensional momenta (normalized by mc) parallel and perpendicular to the mean magnetic field, respectively, σ is an integer ≥ 1 , n_{CR} is the CR number density, v is the CR's velocity written as a function of $u_{\parallel}$ and $u_{\perp}$, $u = (u_{\perp}^2 + u_{\parallel}^2)^{1/2}$, and

$$g(u) = \begin{cases} \frac{p-1}{u_1} \left[1 - \left(\frac{u_1}{u_2} \right)^{p-1} \right]^{-1} \left(\frac{u}{u_1} \right)^{-p}, & u_1 \leq u \leq u_2, \\ 0 & \text{otherwise.} \end{cases} \quad (2)$$

For the standard DSA one has $p = 2$ (Bell 1978; Drury 1983) and when the nonlinear effect on DSA is included, p deviates from this canonical value (Eichler 1984). The distribution with $\sigma = 1$ corresponds to that used in Melrose (1986). The distribution (1) implies

$$\begin{aligned} n_{CR} &= 4\pi \int_0^{\infty} u_{\perp} du_{\perp} \int_{-\infty}^{\infty} du_{\parallel} f(u_{\parallel}, u_{\perp}) \\ &= 2\pi \int_0^{\pi} \sin \alpha d\alpha \int_0^{\infty} u^2 du f(u \cos \alpha, u \sin \alpha), \end{aligned} \quad (3)$$

where $\cos \alpha \equiv u_{\parallel}/u$. In the second expression in (3), one chooses (u, α) in place of $(u_{\parallel}, u_{\perp})$ as independent variables. It can be verified that averaging the parallel velocity $v_{\parallel}$ over the distribution (1) gives the streaming velocity v_{CR} , which is independent of the choice of the parameter σ . The CR current is then given by $J_{CR} = qn_{CR}v_{CR}$. The presence of streaming CRs affects the background plasma in two ways, due to their charge density and their current density, respectively. The background plasma must have a charge density and a current density that are equal and opposite to those of the CRs. This requires that the electrons and ions (assumed to be protons) have different charge densities, $n_e \neq n_p$, and that they move relative to each other with streaming velocities $v_e \neq v_p$, which are assumed to be along the guiding magnetic field. The neutralization conditions require (Achterberg 1983)

$$e(n_e - n_p) = qn_{CR}, \quad e(n_e v_e - n_p v_p) = qn_{CR} v_{CR}. \quad (4)$$

These properties of the background plasma drive the nonresonant instability attributed to the CRs.

2.2 Dispersion relation

A formal procedure to derive the dispersion relation involves separating the plasma response tensor into that for a background plasma, denoted by K_{ij} , plus that for the CR component, denoted by ΔK_{ij} . The background plasma can be regarded as a cold, magnetized plasma, while the CR component is described by the distribution (1). Assume that the gyrofrequency of CRs is $\Omega = |q|B/m$, where B is the mean background magnetic field. A useful approximation is $k_{\perp} v_{\perp} / \tilde{\Omega} \ll 1$, where $k_{\perp}$ is the perpendicular wave number, $\tilde{\Omega} = \Omega/\gamma$ and $\gamma = (1 + u^2)^{1/2}$ is the Lorentz factor of CRs. The approximation implies that in the response tensor, only the first gyroharmonics terms are important. For the background plasma, one assumes $v_A^2 \ll c^2$ and the low-frequency approximations, $\omega \ll \Omega_i$, $\omega \ll |k_{\parallel} v_{\parallel}|$ and $\omega \ll \tilde{\Omega}$, where Ω_i is the gyrofrequency of ions in the background plasma. The background magnetic field is assumed to be along the 3-axis. Since $K_{33} \propto \Omega_i^2/\omega^2$ can be set to ∞ , only the 2×2 components of the response tensor are relevant. For the background plasma, these components can be written as

$$K_{11} = K_{22} = \frac{c^2}{v_A^2}, \quad (5)$$

$$K_{12} = -K_{21} = i \frac{c^2}{v_A^2} \frac{\Omega_i}{\omega} \left[\frac{n_e - n_p}{2n_i} + \frac{k_{\parallel}(n_e v_e - n_p v_p)}{\omega n_i} \right], \quad (6)$$

where $n_{\parallel} = k_{\parallel}c/\omega$, $v_A = \Omega_i/\omega_{pi}$ is the Alfvén speed, ω_{pi} is the ion plasma frequency of the background plasma. Using the neutralization conditions (4), (6) can be rewritten in the form

$$K_{12} = -K_{21} = -in_{\parallel}^2\chi_h^{(0)}, \quad \chi_h^{(0)} = \frac{\mu_0 J_{CR}}{2k_{\parallel}B}. \quad (7)$$

The CR components are

$$\Delta K_{11} = \Delta K_{22} = -in_{\parallel}^2\chi_a, \quad \Delta K_{12} = -\Delta K_{21} = -in_{\parallel}^2\chi_h, \quad (8)$$

where

$$\chi_a = \frac{\pi}{2}A \int_{x_2}^{x_1} dx x^{p+2\sigma-3}(1-x^2)H(1-x^2), \quad (9)$$

$$\chi_h = A \int_{x_2}^{x_1} dx x^{p+2\sigma-3} \left[(1-x^2) \ln \left| \frac{1+x}{1-x} \right| + 2x \right. \\ \left. - 4 \sum_{s=1}^{\sigma-1} \frac{x^{1-2s}}{4s^2-1} \right], \quad (10)$$

$$A = \frac{\mu_0 J_{CR}}{4k_{\parallel}B} \frac{(4\sigma^2-1)(p-1)}{x_1^{p-1}}, \quad (11)$$

with $(x_1, x_2) = b(1/u_1, 1/u_2)$, $b = \Omega/|k_{\parallel}|c$ and $H(x)$ the step function.

Eq (9) and (10) correspond to the dissipative and reactive part respectively of the response tensor due to CRs. In the strong magnetic field limit $x > 1$, the dissipative part is zero. When $x < 1$, both dissipative and reactive terms contribute to the dispersion. The dissipative part is due to CRs in the resonance $x \equiv b/u \approx \mu$ as discussed extensively in the context of the resonant instability (see discussion below) (Kulsrud & Pearce 1969; Melrose & Wentzel 1970; Skilling 1975; McKenzie & Völk 1982; Lagage & Cesarsky 1983a). The dispersion relation takes the following form

$$\omega^2 = \frac{1}{2} \left\{ \frac{k^2}{k_{\parallel}^2} + 1 + 2i\chi_a \right. \\ \left. \pm \left[\frac{k_{\perp}^4}{k_{\parallel}^4} + 4 \left(\chi_h^{(0)} + \chi_h \right)^2 \right]^{1/2} \right\} k_{\parallel}^2 v_A^2. \quad (12)$$

In the $\chi_a \rightarrow 0$ and $\chi_h \rightarrow 0$ limit ($\chi_h^{(0)}$ must also be zero), (12) with the lower sign reproduces the usual Alfvén mode dispersion $\omega = |k_{\parallel}|v_A$ and with the upper sign the fast mode dispersion $\omega = kv_A$.

2.3 Streaming instabilities

Instability occurs when ω has an imaginary part of the appropriate sign. An imaginary part can arise in two different ways, which we refer to as resonant and nonresonant instabilities. Writing $\omega \rightarrow \omega + i\Gamma$ in (12) and assuming $\Gamma \ll \omega$, the resonant instability is described by equating the small imaginary terms, giving $\Gamma = \chi_a k_{\perp}^2 v_A^2 / 2\omega$. The value of χ_a is determined by the CR distribution. A resonant instability can develop for both Alfvén and fast-mode waves if streaming CRs are in resonance with the wave considered. For Alfvén waves ($\omega = |k_{\parallel}|v_A$), the growth rate is

$$\Gamma \approx \frac{1}{2}\chi_a |k_{\parallel}|v_A. \quad (13)$$

The polarization of both modes is generally elliptical and becomes approximately circular for nearly parallel propa-

gation $|k_{\perp}/k_{\parallel}| \ll 1$. In each mode, waves of both sense of polarization can grow.

On neglecting χ_a , (12) with the upper sign becomes a real equation, because the quantity inside the square root is a sum of squares. For the negative sign (Alfvén waves), ω^2 can be negative, and one of the two solutions corresponds to an intrinsically growing wave. This is identified as the nonresonant instability. For $|k_{\perp}/k_{\parallel}| \ll 1$, the condition for this nonresonant instability to occur is $(\chi_h^{(0)} + \chi_h)^2 > 1$. The growth rate is

$$\Gamma \approx \sqrt{2} \left(|\chi_h^{(0)} + \chi_h| - 1 \right)^{1/2} |k_{\parallel}|v_A. \quad (14)$$

In principle, this can be satisfied either due to $|\chi_h| > 1$ or to $|\chi_h^{(0)}| > 1$. For $\sigma = 1$, the integral χ_h given by (10) can be calculated exactly, i.e.,

$$\chi_h = \frac{3\mu_0 J_{CR}(b-1)}{4k_{\parallel}Bx_1^{p-1}} \int_{x_2}^{x_1} dx x^{p-1} \left[(1-x^2) \ln \left| \frac{1+x}{1-x} \right| + 2x \right] \\ = \frac{3\mu_0 J_{CR}}{16k_{\parallel}Bx_1} \left\{ 2 \left[x_1(1+x_1^2) - x_2(1+x_2^2) \right] \right. \\ \left. - (x_1^2-1)^2 \ln \left| \frac{1+x_1}{1-x_1} \right| + (x_2^2-1)^2 \ln \left| \frac{1+x_2}{1-x_2} \right| \right\}. \quad (15)$$

The second equality in (15) is obtained for $p = 2$. The condition $|\chi_h| > 1$ cannot be satisfied for $x_1 \ll 1$. Thus, only the second possibility ($\chi_h^{(0)} > 1$) is relevant.

For $x_2 \ll x_1 \ll 1$, one finds

$$\chi_h \approx \frac{\mu_0 J_{CR}x_1^2}{4k_{\parallel}B}, \quad (16)$$

so that the contribution from the CR current is much smaller than the contribution due to the compensating current in the background plasma. It follows that in treating the nonresonant instability, one can neglect the direct contribution of the CRs, described by χ_h , in comparison with the indirect contribution, described by $\chi_h^{(0)}$. The growth rate (14) can be written in the following approximate form

$$\Gamma \approx \sqrt{2} \left(\chi_h^{(0)} - 1 \right)^{1/2} k_{\parallel}v_A = \sqrt{2} \left(\frac{\mu_0 J_{CR}}{k_{\parallel}B} - 1 \right)^{1/2} k_{\parallel}v_A. \quad (17)$$

Nonresonant instability requires

$$|J_{CR}| > \frac{k_{\parallel}B}{\mu_0}. \quad (18)$$

An explanation for why the direct contribution from the CR current is so much smaller than the indirect contribution from the background plasma is that in the limit $k_{\parallel}r_g \gg 1$, most of CRs move rather rigidly. Their only role in this limit is to induce the compensating current in the background plasma that drives the instability. It is appropriate to point out here that a small fraction of CRs can satisfy the resonant condition ($\mu = x$) and that resonant instability due to these resonant CRs is insignificant compared to the nonresonant instability due to the compensating current. However, the resonant interactions between these CRs and waves with $k_{\parallel} \gg 1/r_g$ can lead to resonant diffusion that has feedback effects on CR streaming motion (cf. Sec. 4).

3 QUASILINEAR DIFFUSION

To determine saturation we calculate the back reaction of wave growth on CRs in the quasilinear diffusion theory (Shapiro & Shevchenko 1964; Davidson 1972). We assume that the nonresonant instability discussed in Sec 2 can develop well into the nonlinear regime, producing turbulence. The growing turbulence causes CRs to diffuse in momentum space and this in turn reduces the anisotropy, suppressing the instability.

We calculate the particle diffusion in the weak turbulence approximation. The distribution of CRs in momentum space can be written as $f(\mathbf{u}, t) = F(u_{\parallel}, u_{\perp}, t) + f^{(1)}(\mathbf{u}, t)$, where $f^{(1)}$ is a linear function of the electric and magnetic fields of the wave and F is the mean distribution, which is assumed to be axisymmetric with respect to the mean magnetic field, $\mathbf{B}$, and as the zeroth order approximation it can be taken to be (1). The response of particles to wave growth is determined by including quadratic terms of the electric and magnetic fields of waves. The diffusion equation for F can be obtained from the Vlasov equation (the derivation is outlined in Appendix). We assume $k_{\perp} r_g \ll 1$ so that only the first gyroharmonics terms are relevant. This condition is generally satisfied for waves propagating nearly parallel to the mean background magnetic field.

3.1 Nonresonant diffusion equation

For a nonresonant instability, it is relevant to consider the approximation that the resonant width is much larger than the growth rate, i.e., $|k_{\parallel} v_{\parallel} - \Omega/\gamma| \gg \Gamma$, where $\Gamma > 0$ is the growth rate and for convenience the frequency is set to zero. The diffusion equation can be expanded in $\Gamma/|k_{\parallel} v_{\parallel} - \Omega/\gamma|$. Let δB_k be the spatially Fourier-transformed magnetic fluctuations arising from the instability. The energy of magnetic fluctuations in an elementary phase volume $d\mathbf{k}/(2\pi)^3$ is $|\delta B_k|^2/2\mu_0$. The expansion yields the following approximate diffusion equation (Appendix A)

$$\begin{aligned} \frac{\partial F}{\partial t} \approx & \frac{1}{2V} \int \frac{d\mathbf{k}}{(2\pi)^3} \Gamma \frac{|\delta B_k|^2}{B_*^2} \left[u_{\perp}^2 \frac{\partial}{\partial u_{\parallel}} \Phi_2 \frac{\partial}{\partial u_{\parallel}} \right. \\ & - (\Phi_1 + \Phi_2) \frac{u_{\parallel}}{u_{\perp}} \frac{\partial}{\partial u_{\perp}} u_{\perp}^2 \frac{\partial}{\partial u_{\parallel}} \\ & - u_{\perp} \frac{\partial}{\partial u_{\parallel}} (\Phi_2 - \Phi_1) u_{\parallel} \frac{\partial}{\partial u_{\perp}} \\ & \left. + \Phi_2 \frac{u_{\parallel}^2}{u_{\perp}} \frac{\partial}{\partial u_{\perp}} u_{\perp} \frac{\partial}{\partial u_{\perp}} \right] F, \end{aligned} \quad (19)$$

$$\Phi_1 = \frac{1}{u_{\parallel}^2 - b^2}, \quad \Phi_2 = \frac{u_{\parallel}^2 + b^2}{(u_{\parallel}^2 - b^2)^2}, \quad (20)$$

with $b = B/B_*$, $B_* = |k_{\parallel}|mc/|q|$ and V the volume of the region considered.

3.2 Perpendicular and parallel kinetic energy

It is of interest to examine the evolution of the parallel and perpendicular kinetic energy by evaluating the rate of change in the average $u_{\parallel}^2$ and $u_{\perp}^2$ (over the mean distribution F). The averaging involves singular terms $1/(u_{\parallel} \pm b)^n$ with

$n = 1, 2, 3$. The $n \geq 2$ terms can be avoided using

$$\Phi_2 = -\Phi_1 + \frac{u_{\parallel}^2}{b} \frac{\partial \Phi_1}{\partial b}, \quad (21)$$

$$\frac{\partial \Phi_1}{\partial u_{\parallel}} = -\frac{u_{\parallel}}{b} \frac{\partial \Phi_1}{\partial b}, \quad (22)$$

$$\frac{\partial \Phi_2}{\partial u_{\parallel}} = \frac{3u_{\parallel}}{b} \frac{\partial \Phi_1}{\partial b} - \frac{u_{\parallel}^3}{b} \frac{\partial}{\partial b} \frac{1}{b} \frac{\partial \Phi_1}{\partial b}. \quad (23)$$

Multiplying (19) by $u_{\parallel}^2$ and $u_{\perp}^2$, respectively, and integrating them over $2\pi u_{\perp} du_{\perp} du_{\parallel}$ divided by n_{CR} , one obtains

$$\begin{aligned} \frac{d\langle u_{\parallel}^2 \rangle}{dt} = & \frac{1}{2\tau} \left[-\langle (u_{\perp}^2 - 4u_{\parallel}^2) \Phi_1 \rangle + \frac{2}{b} \frac{\partial}{\partial b} \langle (2u_{\perp}^2 - u_{\parallel}^2) u_{\parallel}^2 \Phi_1 \rangle \right. \\ & \left. - \frac{1}{b} \frac{\partial}{\partial b} \frac{1}{b} \frac{\partial}{\partial b} \langle u_{\parallel}^4 \Phi_1 u_{\perp}^2 \rangle \right], \end{aligned} \quad (24)$$

$$\begin{aligned} \frac{d\langle u_{\perp}^2 \rangle}{dt} = & \frac{1}{2\tau} \left[-2\langle \Phi_1 u_{\parallel}^2 \rangle + \frac{1}{b} \frac{\partial}{\partial b} \langle u_{\parallel}^2 \Phi_1 (2u_{\parallel}^2 - 3u_{\perp}^2) \rangle \right. \\ & \left. + \frac{1}{b} \frac{\partial}{\partial b} \frac{1}{b} \frac{\partial}{\partial b} \langle u_{\parallel}^4 \Phi_1 u_{\perp}^2 \rangle \right], \end{aligned} \quad (25)$$

where

$$\tau = \left(\frac{1}{V} \int \frac{d\mathbf{k}}{(2\pi)^3} 2\Gamma \frac{|\delta B_k|^2}{B_*^2} \right)^{-1}. \quad (26)$$

The ratio of the parallel to perpendicular pressures is determined by the same angular averages as the ratio of $\langle u_{\parallel}^2 \rangle$ to $\langle u_{\perp}^2/2 \rangle$. It follows that the ratio of Eqs (24) and (25) effectively determines how the ratio of the parallel to perpendicular pressures evolves.

3.3 Long wavelength approximation

In the long wavelength approximation $1/k_{\parallel} \gg r_g$, one has $b \gg |u_{\parallel}|$ and

$$\Phi_1 \approx -\Phi_2 \approx -\frac{1}{b^2}. \quad (27)$$

The rates of change in the average of $u_{\parallel}^2$ and $u_{\perp}^2$ is given by

$$\frac{d\langle u_{\parallel}^2 \rangle}{dt} = -\left(2\langle u_{\parallel}^2 \rangle - \langle u_{\perp}^2 \rangle \right) \frac{1}{\tau'}, \quad (28)$$

$$\frac{d\langle u_{\perp}^2 \rangle}{dt} = \langle u_{\parallel}^2 \rangle \frac{1}{\tau'}, \quad (29)$$

where τ' has the same form as (26) but with B_* replaced by B . Thus, one has $1/\tau' = \partial(\delta B^2/B^2)/\partial t$. One may compare the nonresonant diffusion to the firehose instability. Similar to the firehose instability, a growing wave causes the parallel kinetic energy to decrease and the perpendicular energy to increase with time. For the CR streaming distribution (1), one can show that the right-hand sides of (28) and (29) are zero. In the long wavelength regime an instability can arise from both dissipative (resonant) and reactive (nonresonant) effects. As we focus on the nonresonant instability in the short wavelength regime, quasilinear diffusion in the long wavelength regime is not discussed further.

3.4 Short wavelength approximation

The right-hand side of Eq (24) and (25) can be evaluated in a general case in which the particle distribution is isotropic in pitch angles. The angular integration in (24) and (25) leads to

$$\frac{d\langle u_{\parallel}^2 \rangle}{dt} = \frac{1}{\tau} \left\langle 2 + x \ln \left| \frac{1-x}{1+x} \right| \right\rangle, \quad (30)$$

$$\frac{d\langle u_{\perp}^2 \rangle}{dt} = -\frac{1}{\tau} \left\langle \frac{2}{3} + 2x^2 + x^3 \ln \left| \frac{1-x}{1+x} \right| \right\rangle, \quad (31)$$

where one uses the following integral

$$\begin{aligned} I_{2n} &\equiv u^2 \int_{-1}^1 \mu^{2n} \Phi_1 d\mu \\ &= 2 \sum_{s=1}^n \frac{x^{2(n-s)}}{2s-1} + x^{2n-1} \ln \left| \frac{1-x}{1+x} \right|. \end{aligned} \quad (32)$$

The second expression in (32) is obtained by retaining the Cauchy principal value. Surprisingly, Eq (30) and (31) have an opposite sign compared to their counterpart in the long wavelength regime (Eq 28 and 29). Although both the firehose instability and the instability discussed here are driven by anisotropy in the particle distribution in momentum space, there is an important difference between the two. The firehose instability is driven by anisotropy in kinetic energy with $\langle u_{\parallel}^2 \rangle > \langle u_{\perp}^2 \rangle/2$. By contrast, the streaming instability is due to the CR streaming motion ($\langle u_{\parallel} \rangle \neq 0$), which is also called streaming anisotropy, and the instability can develop even when the CR pressure is isotropic. With the specific choice of the distribution (1), one can show that $\langle u_{\parallel}^2 \rangle = \langle u_{\perp}^2 \rangle/2$. Since the current in (1) is $\propto \mu^{2\sigma-1}$, it does not contribute to the average $\langle Y \rangle$ if Y is proportional to an even power of μ .

The instability can lead to pressure anisotropy with perpendicular pressure increasing. In the approximation $x \ll 1$, one has $d\langle u_{\parallel}^2 \rangle/dt \approx 2/\tau$ and $d\langle u_{\perp}^2 \rangle/dt \approx -2/3\tau$. This gives

$$\frac{d}{dt}(2\langle u_{\parallel}^2 \rangle - \langle u_{\perp}^2 \rangle) = \frac{10}{3}\tau^{-1} > 0. \quad (33)$$

One should emphasize here that development of such anisotropy in kinetic energy is driven by the streaming motion of CRs.

From (19), one may calculate the rate of change in the streaming speed due to nonresonant diffusion. One starts with a particle distribution with streaming motion similar to (1) but without specifying the specific form of $g(p)$. Consider the case $\sigma = 1$, i.e., the streaming component is $\propto \mu$. As in Sec 4.4, in the averaging, one can carry out the angular integration first. This gives

$$\begin{aligned} \frac{d\langle v_{\parallel} \rangle}{dt} &\approx -\frac{1}{4\tau} \left\langle \frac{v}{u^2} \left[-(4 + \beta^2)I_2 - (2 - 8\beta^2 + 3\beta^4)I_4 \right. \right. \\ &\quad \left. \left. - 3\beta^2(1 - \beta^2)I_6 + \frac{1}{x} \frac{\partial}{\partial x} \left[-3I_2 + (1 + 5\beta^2)I_4 \right. \right. \right. \\ &\quad \left. \left. \left. + \beta^2(6 - 7\beta^2\beta^2)I_6 + 3\beta^4I_8 \right] \right. \right. \\ &\quad \left. \left. + \frac{1}{x} \frac{\partial}{\partial x} \frac{1}{x} \frac{\partial}{\partial x} \left[I_4 - \beta^2I_6 - (1 - \beta^2)I_8 \right] \right] \right\rangle, \end{aligned} \quad (34)$$

where I_{2n} is given by (32). Since $dv_{\text{CR}}/dt = d\langle v_{\parallel} \rangle/dt$, in the limit $x \ll 1$, one has

$$\frac{dv_{\text{CR}}}{dt} \approx -\frac{3v_{\text{CR}}}{\tau} \frac{2}{5} \left\langle \frac{1}{u^2} (-70 + 28\beta^2 + 3\beta^4) \right\rangle. \quad (35)$$

One obtains $dv_{\text{CR}}/dt > 0$, contrary to what one would expect. In Sec 4, cf. Eq (40), we show that that resonant diffusion has dominant effects on the evolution of CR streaming motion, which ensures $dv_{\text{CR}}/dt < 0$.

4 RESONANT DIFFUSION

Waves generated from the nonresonant CR-streaming instability can interact with CRs in resonance at $\mu = x$. Although the fraction of CRs in the resonance is small, we show that diffusion through such resonant interactions has a dominant effect on the CR streaming motion. Provided that the phase speed of the wave is $\ll c\mu$, resonant diffusion is equivalent to pitch angle scattering, in which particles change their direction of motion without gaining or losing energy. Since the nonresonant instability is in the regime $x \ll 1$, resonant scattering involves mainly CRs with pitch angles $\alpha \sim \pi/2 \pm x$. Since one deals with purely growing waves that can be regarded as a quasi-mode with $\omega \rightarrow 0$, the assumption of low phase speed $\ll c\mu$ is valid even for particles with pitch angles near $\pi/2$. The standard formalism involves writing down the diffusion equation involving pitch angles only (Melrose 1986). Using $F(u, \mu) = F(u_{\parallel}, u_{\perp})$ with the replacements $u_{\parallel} = u\mu$ and $u_{\perp} = u\sqrt{1 - \mu^2}$ and retaining only the pitch angle scattering part in (A12), one obtains

$$\begin{aligned} \frac{dF(u, \mu)}{dt} &= \frac{\partial}{\partial \mu} \left[(1 - \mu^2) D \frac{\partial}{\partial \mu} \right] F(u, \mu), \\ D &= \frac{\pi}{4} \frac{\Omega}{\gamma} \frac{\delta B_r^2}{B^2}, \end{aligned} \quad (36)$$

where

$$\frac{\delta B_r^2}{\mu_0} = \frac{k_r}{V} \int \frac{dk_{\perp} k_{\perp}}{(2\pi)^2} \frac{\delta B_k^2}{\mu_0} \Big|_{k_r} \quad (37)$$

is the density of magnetic energy at the resonant wavenumber $k_r = \Omega/cu|\mu|$. Using (1), it can be shown that the resonant diffusion does not contribute to change in $\langle u_{\parallel}^2 \rangle$ and $\langle u_{\perp}^2 \rangle$. Thus, the rate of change in parallel and perpendicular pressures is determined by nonresonant diffusion.

The rate of change in the streaming speed is

$$\frac{dv_{\text{CR}}}{dt} = -\bar{D}v_{\text{CR}}, \quad (38)$$

where

$$\bar{D} = \frac{1}{2} \int_{-1}^1 d\mu (1 - \mu^2) D \approx \frac{\pi^2}{4} \frac{\Omega}{\gamma} \frac{\delta B_r^2}{B^2} \frac{1}{k_0 r_{g0}}, \quad (39)$$

where δB_r is set the value at $\mu \sim \pi/2$. The approximation in (39) is obtained by writing $\delta B_k^2 = (2\pi)^3 V \delta B_r^2 \delta(k_{\perp}) \delta(k_r - k_0)/k_{\perp}$, i.e. one assumes that the magnetic fluctuations are centered at the wave number k_0 . Here r_{g0} is the gyroradius of CRs with an average Lorentz factor $\gamma_0 = \langle \gamma \rangle$.

Since the rate of the change in streaming motion due to resonant diffusion is $(dv_{\text{CR}}/dt)_r \propto \Omega/\gamma$ and the correspond-

ing rate due to nonresonant diffusion is $(dv_{\text{CR}}/dt)_{nr} \propto \Gamma/\gamma^2$, one has

$$\left| \frac{(dv_{\text{CR}}/dt)_r}{(dv_{\text{CR}}/dt)_{nr}} \right| \sim 5 \times 10^{-2} \frac{\gamma\Omega}{\Gamma} \left(\frac{\delta B_r^2}{\delta B^2} \right) \left(\frac{B_*^2}{B^2} \right) \frac{1}{k_0 r_{g0}}. \quad (40)$$

Here δB is the typical amplitude of the magnetic fluctuations generated by the nonresonant instability. The right-hand side is larger than 1 for $k_0 r_{g0} > 20(\Gamma/\gamma_0\Omega)$, where one assumes $\delta B_r \sim \delta B$ and $B_*/B \sim k_0 r_{g0}$. This condition is satisfied for the typical parameters relevant for SN shocks (cf. Sec 5). If one writes $\delta B \propto \exp(\Gamma t)$ and takes the limit $t \gg 1/\Gamma$, Eq (38) can be solved to yield

$$\frac{\delta B^2}{B^2} \approx \frac{12}{\pi} \frac{\Gamma\gamma}{\Omega} k_0 r_{g0} \ln \left(\frac{v_{\text{CR}0}}{v_{\text{CR}}} \right). \quad (41)$$

5 MAGNETIC FIELDS IN SNRS

There are a number of SNRs whose post-shock magnetic fields have been estimated from X-ray observations and found to be in the range $\geq 10^{-8}$ T, much higher than the typical value in the ISM ($\sim 10^{-10}$ T) (Vink & Laming 2003; Völk, Berezhko & Ksenofontov 2005). Although one cannot rule out the possibility that such strong field is due to compression of strong pre-existing magnetic fields in the stellar wind of the progenitor star, the observations favor the interpretation that the magnetic fields are due to an instability at the shock. Observations show that the magnetic field strength strongly depends on the shock speed (Vink & Laming 2003; Völk, Berezhko & Ksenofontov 2005). The post-shock magnetic energy density in SNRs whose shock speeds have been estimated show a tendency $\delta B^2 \propto v_s^\delta$, where v_s is the shock speed (Völk, Berezhko & Ksenofontov 2005; Vink 2008). The argument that the magnetic pressure is proportional to the ram pressure ($\propto v_s^2$) at the shock front suggests $\delta = 2$, while Bell (2004)'s model predicts $\delta = 3$. So far, current observations cannot differentiate these two because of the narrow dynamic range of the observed properties of SN shocks (Vink 2008).

5.1 SN shocks

We consider a forward shock traveling at a velocity v_s . The shock undergoes initially a free expansion phase and after sweeping up sufficient mass, it enters a Sedov phase. In the Sedov phase, the velocity is given by $v_s = 2R_s/5t$, where R_s is the shock radius. The total kinetic energy of the shock is $E = (4/3)R_s^3 \rho_0 v_s^2$, where ρ_0 is the density of the ISM. On eliminating R_s in favour of t , the shock velocity becomes

$$v_s = \left(\frac{6E}{125\rho_0} \right)^{1/5} t^{-3/5}. \quad (42)$$

One can show that development of the nonresonant instability is faster in the early Sedov phase. The maximum growth rate is $\propto J_{\text{CR}}^{1/2}$, which can be written in terms of the ram pressure in the up-stream region. Thus, one expects the growth rate to be a strong function of the shock speed. Since the ram pressure is $\sim \rho_0 v_s^2$, the CR pressure can be expressed as $P_{\text{CR}} = \eta_s \rho_0 v_s^2$, where $\eta_s \leq 1$ is the acceleration efficiency.

We consider precursor CRs with the initial streaming velocity $v_{\text{CR}0} \sim v_s$. Since $P_{\text{CR}} = n_{\text{CR}} mc \langle uv \rangle / 3$, one has $J_{\text{CR}} = 3\eta_s q \rho_0 v_s^2 v_{\text{CR}} / (mc \langle uv \rangle) \approx 3\eta_s q \rho_0 v_s^2 v_{\text{CR}} / [mc u_1 \ln(u_2/u_1)]$, where $\langle uv \rangle \approx cu_0$ and u_0 is the average momentum given by

$$u_0 = \int_0^\infty u g(u) du = u_1 \frac{p-1}{p-2} \frac{1 - (u_2/u_1)^{2-p}}{1 - (u_2/u_1)^{1-p}}. \quad (43)$$

The average Lorentz factor of CRs is $\gamma_0 = (1 + u_0^2)^{1/2}$. For $p = 2$ and $u_2 \gg u_1$, one has $u_0 = u_1 \ln(u_2/u_1)$. Therefore, the faster the shock speed the higher the growth rate.

5.2 Saturated magnetic field vs shock speed

One may estimate the maximum magnetic field, δB_m , of the wave by equating the magnetic pressure to the CR pressure, $\delta B_m^2 / 2\mu_0 = P_{\text{CR}}$. This leads to $\delta B_m \approx (2\mu_0 \eta_s \rho_0)^{1/2} v_s$, or

$$\delta B_m \approx 3 \times 10^{-7} \eta_s^{1/2} \left(\frac{n_0}{10^6 \text{ m}^{-3}} \right)^{1/2} \left(\frac{v_s}{5 \times 10^6 \text{ m s}^{-1}} \right) \text{ T}. \quad (44)$$

Eq (44) is a rather optimistic estimate and calculation of the saturation due to the feedback effects of the instability on the streaming anisotropy generally leads to a lower saturation level than (44).

One can calculate the saturated magnetic field due to the diminishing of the streaming anisotropy from (41). This involves calculation of the cut-off streaming speed, denoted by v'_{CR} , at which the instability turns off. From (18), one obtains

$$v'_{\text{CR}} \approx \frac{5c}{24\eta_s} \frac{U_B}{\rho_0 v_s^2} k_{\parallel} r_{g0}, \quad (45)$$

where $U_B = B^2 / 2\mu_0$ is the energy density of the mean magnetic field. At a given parallel wavenumber $k_{\parallel}$, wave growth stops when the streaming speed is reduced to $v_{\text{CR}} \leq v'_{\text{CR}}$. One may assume $v_{\text{CR}0} \sim v_s$. Eq (45) gives

$$\frac{v_{\text{CR}0}}{v'_{\text{CR}}} \approx 5 \times 10^5 \frac{\eta_s}{k_{\parallel} r_{g0}} \left(\frac{v_s}{5 \times 10^6 \text{ m s}^{-1}} \right)^3 \left(\frac{10^{-10} \text{ T}}{B} \right)^2 \times \left(\frac{n_0}{10^6 \text{ m}^{-3}} \right). \quad (46)$$

For convenience, one assumes here that CRs and ions in the ISM are all protons.

The growth rate (17) is estimated to be $\Gamma \approx (6\eta_s k_0 r_{g0})^{1/2} (v_s/c)^{3/2} \Omega / \gamma_0$. From (41) one obtains

$$\frac{\delta B^2}{B^2} \approx \frac{12}{\pi} (6\eta_s)^{1/2} \left(\frac{v_s}{c} \right)^{3/2} (k_0 r_{g0})^{3/2} \ln \left(\frac{v_{\text{CR}0}}{v'_{\text{CR}}} \right). \quad (47)$$

Eq (47) implies that significant amplification $\delta B^2/B^2 \gg 1$ is possible if one assumes a large $k_0 r_{g0}$. However, one should point out that the quasilinear theory is based on the weak turbulence assumption that requires the relevant wave fluctuations be small. Nonetheless, since the saturation mechanism discussed is quite generic, i.e. the saturation is attributed to the reduction in CR streaming, one expects that extrapolation of Eq (47) to the $\delta B^2/B^2 \gg 1$ regime would still provide a reasonable estimate for the saturation. To estimate the upper limit to the saturation one assumes that the Larmor radius is limited by the scale length of the acceleration region, which is assumed to be the shock's radius, R_s . The characteristic wavenumber k_0 is limited by the condition (18), which gives $k_0 \sim (\omega_{p,\text{CR}}/c)(v_{\text{CR}}/v_A)$, where

$\omega_{p,CR}$ is the plasma frequency of CRs. This corresponds to $(k_0 r_{g0})_{\max} \sim 10^2 - 10^3$. For $\eta_s = 0.1$, $v_s = 5 \times 10^6 \text{ m s}^{-1}$ and $k_0 r_{g0} = 10^2$, one has $\delta B^2/B^2 \sim 39$. The saturated magnetic field can exceed the background field by a modest factor. If the CR pressure is a fixed fraction of the ram pressure, one has $n_{CR} \gamma_0 \propto v_s^2$. Since the shock radius can be expressed as $R_s = (3E/4\rho_0 v_s^2)^{1/3}$, one has $k_0 R_s \propto v_s^{1/3}$, where $k_0 \propto v_{CR} \sim v_s$. Eq (47) implies that the energy density of the saturated magnetic field increases with the shock speed as $\delta B^2 \propto v_s^2$. Remarkably, this predicts the same shock-speed scaling as the one from the equipartition argument, i.e. the magnetic pressure equal to the CR pressure (cf. Eq 44).

6 CONCLUSIONS AND DISCUSSION

We develop a kinetic version of the CR streaming instability of Bell (2004), who treated it using MHD. We show that the nonresonant growth is not due to the CRs themselves, but rather to a current in the background plasma that neutralizes the current due to streaming CRs. Saturation of the CR streaming instability is discussed in the quasilinear theory. Growth of the nonresonant instability leads to particle diffusion that suppresses the instability. In the quasilinear diffusion formalism, the saturated magnetic field can be calculated analytically. We consider both resonant and nonresonant diffusion, with the latter being treated in a similar way to that used for the firehose instability (Shapiro & Shevchenko 1964; Davidson 1972). In our model the CR pressure is set to a fixed fraction of the ram pressure at the shock front. It is shown that evolution of the parallel and perpendicular pressures is due to nonresonant diffusion. The direction of the evolution is that the parallel energy increases and perpendicular energy decreases. This is in contrast to the firehose instability which causes the perpendicular pressure to increase and parallel pressure to decrease. One should note the difference in the sources of free energy that drive the instabilities; the free energy for the firehose instability is an excess of parallel pressure while the free energy for the nonresonant instability considered here is streaming motion of CRs. It is shown that saturation is determined by resonant diffusion which reduces the streaming motion. The saturated magnetic fluctuations can exceed the background field provided that the shocks are sufficiently fast and saturation occurs at short wavelength $1/k_0 \ll 1$.

There have been numerical simulations of the streaming instability in both the MHD approximation and kinetic theory, which confirm rapid growth of the instability under the physical conditions of young SN shocks. However, these simulations predict different saturation levels. One of the main difficulties in numerical simulation of the instability is the huge difference between the CR number density and background plasma density, with $n_{CR}/n_0 \sim 10^{-5}$. A particular approximation is usually adopted to make numerical simulation of such system feasible. For example, in Niemiec, Pohl, Stroman, & Nishikawa (2008)'s numerical model, the density ratio is set artificially to a large value. Because the existing numerical models use different approximations, and it is unrealistic to make meaningful comparison among their results. MHD simulations generally predict a saturation at strong mag-

netic fields (Bell 2004; Zirakashvili, Ptuskin & Völk 2008), while kinetic particle-in-cell (PIC) simulations appear to predict much lower saturated magnetic fields (Niemiec, Pohl, Stroman, & Nishikawa 2008; Riquelme & Spitkovsky 2008). The analytical calculation presented here predicts a relatively modest amplification that gives rise to magnetic field fluctuations similar to the level predicted by the recent kinetic simulations (Riquelme & Spitkovsky 2008).

REFERENCES

- Achterberg, A., 1983, *A&A*, 119, 274
- Amato, E., Blasi, P., 2008, *MNRAS*, in press
- Bell, A. R., 1978, *MNRAS*, 182, 147
- Bell, A. R., 2004, *MNRAS*, 353, 550
- Bell, A. R., Lucek, S. G., 2001, *MNRAS*, 321, 433
- Davidson, R. C., 1972, *Methods in Nonlinear Plasma Theory* (Academic Press: New York)
- Drury, L. O'C., 1983, *Rep. Prog. Phys.*, 46, 973
- Eichler, D., 1984, *ApJ*, 277, 429
- Hillas, A. M., 2006, *J. Phys. Conf. Series* 47, 168
- Kulsrud, R., Pearce, W. P., 1969, *ApJ*, 156, 445
- Lagage, P. O., Cesarsky, C. J., 1983a, *A&A*, 118, 223
- Lagage, P. O., Cesarsky, C. J., 1983b, *A&A*, 125, 249
- Machabeli, G. Z., Luo, Q., Melrose, D. B., Vladimirov, S., 2000, *MNRAS*, 312, 51
- McKenzie, J. F., Völk, H. J., 1982, *A&A*, 116, 191
- Melrose, D. B., Wentzel, D. G., 1970, *ApJ*, 457, 476
- Melrose, D. B., 2005, *AIP Proceedings*, 781, 135
- Melrose, D. B., 1986, *Instabilities in Space and Laboratory Plasmas*, Cambridge Press
- Niemiec, J., Pohl, M., Stroman, T., Nishikawa, K. I., 2008, *ApJ*, 684, 1174
- Reville, B., Kirk, J. G., Duffy, P., 2006, *Plasma Phys. Control. Fusion*, 48, 1741
- Riquelme, M. A., Spitkovsky, A., 2008, *ApJ*, submitted
- Shapiro, V. D., Shevchenko, V. I., 1964, *Sov. Phys. JETP* 18, 1109
- Skilling, J., 1975, *MNRAS*, 172, 557
- Tanaka, T., et al., 2008, *ApJ*, in press
- Vink, J., 2008, *astro-ph* 0810.3680
- Vink, J., Laming, J. M., 2003, *ApJ*, 584, 758
- Völk, H., Berezhko, E. G., Ksenofontov, L. T., 2005, *A&A*, 433, 229
- Völk, H., Ksenofontov, L. T., Berezhko, E. G., 2008, *astro-ph*0809.2432
- McKenzie, J. F., Völk, H., 1982, 17th Int. Cosmic Ray Conf., Vol. 9, 242
- Weibel, E. S., *Phys. Rev. Lett.* 2, 83
- Zirakashvili, V. N., Ptuskin, V. S., Völk, H. J., 2008, *ApJ*, 678, 255

APPENDIX A: QUASILINEAR DIFFUSION

Using the similar method discussed in Davidson (1972), we write the distribution function as $f \approx f^{(0)} + f^{(1)}$ with $f^{(1)}$ a linear function of electric and magnetic fluctuations, $\delta \mathbf{E}$ and $\delta \mathbf{B}$, of the wave. A formal approach to determine the

feedback effect of the wave on the particle distribution involves calculation of time-dependence of the distribution averaged over the random phase, denoted by $F \equiv \langle f^{(0)} \rangle_R$, where $\langle \cdots \rangle_R$ is the random phase average.

A1 General formalism

From the Vlasov equation one obtains

$$\frac{dF}{dt} + q \left\langle (\delta \mathbf{E} + \mathbf{v} \times \delta \mathbf{B}) \cdot \frac{\partial f^{(1)}}{\partial \mathbf{p}} \right\rangle_R = 0, \quad (\text{A1})$$

where $d/dt = \partial/\partial t + \mathbf{v}_D \cdot \partial/\partial \mathbf{x}$, $\mathbf{v}_D$ is the drift velocity. Let $\xi \equiv (\delta \mathbf{E}, \delta \mathbf{B}, f^{(1)})$ and the corresponding spatial Fourier transform be $\xi_k \equiv (\delta \mathbf{E}_k, \delta \mathbf{B}_k, f_k^{(1)})$, defined by

$$\xi(\mathbf{x}, t) = \int \frac{d\mathbf{k}}{(2\pi)^3} \xi_k(t) \exp(i\mathbf{k} \cdot \mathbf{x}), \quad (\text{A2})$$

where we assume WKB for the time dependence

$$\xi_k(t) \propto \exp[-i\psi_k(t)], \quad \psi_k(t) = \int_0^t [\omega_k(t) + i\Gamma(t)] dt'. \quad (\text{A3})$$

Here ω_k is the wave frequency and Γ is the growth rate. One adopts the convention $\omega_{-k} = -\omega_k$, $\xi_{-k} = \xi_k^*$.

In cylindrical coordinates, the velocity can be expressed as $\mathbf{v} = (v_\perp \cos \phi, v_\perp \sin \phi, v_\parallel)$, where ϕ is the gyrophase. The 3-axis is assumed to be along the magnetic field. For a transverse wave propagating parallel to the magnetic field, one has $\delta E_\pm \equiv \delta E_{kx} \pm \delta E_{ky}/i = \mp(i\omega/k_\parallel)(\delta B_{kx} \pm \delta B_{ky}/i) \equiv \mp(i\omega/k_\parallel)\delta B_\pm$. In the limit $k_\perp v_\perp/\Omega \ll 1$, the perturbation of the distribution can be written as

$$f_k^{(1)} = \frac{1}{2} \left(f_+ e^{i\phi} + f_- e^{-i\phi} \right). \quad (\text{A4})$$

From the linearized Vlasov equation one obtains

$$f_\pm = \frac{\pm q \delta B_\pm}{\omega + i\Gamma - k_\parallel v_\parallel \pm \Omega/\gamma} \times \left[v_\perp \frac{\partial}{\partial p_\parallel} - \left(v_\parallel - \frac{\omega + i\Gamma}{k_\parallel} \right) \frac{\partial}{\partial p_\perp} \right] F. \quad (\text{A5})$$

Substituting (A4) and (A5) into (A1), one derives

$$\begin{aligned} \frac{dF}{dt} = & -\frac{q}{4iV} \int \frac{d\mathbf{k}}{(2\pi)^3} \left[v_\perp \frac{\partial}{\partial p_\parallel} \right. \\ & \left. - \left(v_\parallel - \frac{\omega - i\Gamma}{k_\parallel} \right) \frac{1}{p_\perp} \frac{\partial}{\partial p_\perp} p_\perp \right] \\ & \times (\delta B_+^* f_+ - \delta B_-^* f_-), \end{aligned} \quad (\text{A6})$$

where one uses

$$\langle \delta \mathbf{B}_k \cdot \delta \mathbf{B}_{k'} \rangle_R = \frac{(2\pi)^3}{V} |\delta B_k|^2 \delta(\mathbf{k} + \mathbf{k}'). \quad (\text{A7})$$

With $|\delta B_k|^2 = |\delta B_+|^2 = |\delta B_-|^2$, one obtains

$$\begin{aligned} \frac{dF}{dt} = & -\frac{q}{2iV} \int \frac{d\mathbf{k}}{(2\pi)^3} |\delta B_k|^2 \\ & \times \left[v_\perp \frac{\partial}{\partial p_\parallel} - \left(v_\parallel - \frac{\omega - i\Gamma}{k_\parallel} \right) \frac{1}{p_\perp} \frac{\partial}{\partial p_\perp} p_\perp \right] \end{aligned}$$

$$\begin{aligned} & \times \frac{\omega + i\Gamma - k_\parallel v_\parallel}{(\omega + i\Gamma - k_\parallel v_\parallel)^2 - \Omega^2/\gamma^2} \\ & \times \left[v_\perp \frac{\partial}{\partial p_\parallel} - \left(v_\parallel - \frac{\omega + i\Gamma}{k_\parallel} \right) \frac{\partial}{\partial p_\perp} \right] F. \end{aligned} \quad (\text{A8})$$

For practical purposes, we set $\omega \sim 0$ here.

A2 Nonresonant diffusion

The quasilinear diffusion equation in the long wavelength approximation can be obtained using the following expansion in $1/\Omega$:

$$\frac{\omega + i\Gamma - k_\parallel v_\parallel}{(\omega + i\Gamma - k_\parallel v_\parallel)^2 - \Omega^2/\gamma^2} \approx \frac{k_\parallel v_\parallel}{\tilde{\Omega}^2} - \frac{i\Gamma}{\tilde{\Omega}^2}, \quad (\text{A9})$$

with $\tilde{\Omega} = \Omega/\gamma$. We consider a more general expansion in $\Gamma/|k_\parallel v_\parallel - \tilde{\Omega}| \gg 1$:

$$\begin{aligned} \frac{i\Gamma - k_\parallel v_\parallel}{(i\Gamma - k_\parallel v_\parallel)^2 - \tilde{\Omega}^2} & \approx \frac{1}{k_\parallel^2 v_\parallel^2 - \tilde{\Omega}^2} \\ & \times \left(i\Gamma - k_\parallel v_\parallel - \frac{2i\Gamma k_\parallel^2 v_\parallel^2}{k_\parallel^2 v_\parallel^2 - \tilde{\Omega}^2} \right). \end{aligned} \quad (\text{A10})$$

Integration on the right-hand side of (A8) includes terms $\propto k_\parallel$ or $\propto 1/k_\parallel$. For turbulent spectra with symmetry $k_\parallel \rightarrow -k_\parallel$, one has

$$\int_{-\infty}^{\infty} dk_\parallel \left(\text{terms} \propto k_\parallel \text{ or } \propto 1/k_\parallel \right) = 0, \quad (\text{A11})$$

and only terms $\propto i\Gamma$ remain.

A3 Resonant diffusion

The standard procedure to write down the resonant diffusion involves setting $\Gamma = 0$. Eq (A8) can be expressed in the following familiar form

$$\begin{aligned} \frac{dF}{dt} = & \frac{\partial}{\partial p_\parallel} \left(D_{\parallel\parallel} \frac{\partial}{\partial p_\parallel} + D_{\parallel\perp} \frac{\partial}{\partial p_\perp} \right) \\ & + \frac{1}{p_\perp} \frac{\partial}{\partial p_\perp} \left(D_{\perp\parallel} \frac{\partial}{\partial p_\parallel} + D_{\perp\perp} \frac{\partial}{\partial p_\perp} \right), \end{aligned} \quad (\text{A12})$$

where $D_{\alpha\beta}$ are the diffusion coefficients.